PRIME S

UNVEILING OUR COSMIC ORIGINS IN THE FAR INFRARED

Jesan Dhi

Dr. Jason Glenn, Principal Investigator Goddard Space Flight Center and Jet Propulsion Laboratory


Jan 9, 2024 Matt Bradford (Jet Propulsion Laboratory, California Institute of Technology) on behalf of the Jason and the PRIMA

team.



PI science overview Instrumentation Detectors

GO science



Pre-decisional information – for planning and discussion purposes only.

PRIMA at a Glance

- 1.8-m, all-aluminum telescope cooled to 4.5 K.
- PRIMAger imager and polarimeter (France / Holland): 25-80 microns R=10 hyperspectral imaging, 91-232 μm imaging polarimetry.

Ó

Ø

Ø

Ø


Improvement is comparable to day

/ night difference in the optical.

Requires a cold (~5K) telescope


and sensitive far-IR detectors.

- FIRESS Spectrometer (JPL w/ GSFC) : 24-235 μm in 4 grating modules with R>85. High-res mode gives R of thousands across full band.
- 100 mK focal planes with kinetic inductance detectors, provided by joint JPL/ GSFC and SRON team.
- JPL lead with GSFC, Ball spacecraft, IPAC data handling, many others.




PR'MA

PRIMA PI Science Programs: Exercises the observatory, and provides legacy datasets



3

PRIMAger Instrument (Leads Denis Burgarella, Laure Ciesla, Marc Sauvage)




- 2 hyperspectral focal planes using linear variable filters with continuous R=10 coverage.
- 4 single-band polarimetric focal planes
- Whole instrument read out simultaneously.


Expect ~ 10,000 galaxies per delta z = 0.25 bin. A

rich archival dataset.

Source Confusion with PRIMA




- Nearly all PRIMA sources will have position and redshift priors with Euclid / Roman surveys.
- Beamsize with PRIMA at 55-60 µm will be comparable to Spitzer 24 µm, which was unconfused and resolved most of the light. Expect the same for PRIMA but reaching further into the far-IR. Use shortwavelengths positions as priors to long-wave extraction (see right).
- Black hole growth / star formation ratio is measured with fluxes shortward of 90 microns.
- Confusion mitigated with spectroscopy – unconfused in line counts.




- XID+ deblending demonstrated with Herschel PACs and SPIRE datasets.
- Uses prior information on both position and flux.
 - Starts with shortest-wavelength map, uses previous flux as prior.
 - CIGALE-based approach uses prior based on all shorter-wavelength data. -> 8x improvement in

depth

Simulated PRIMAger hyperspectral dataset



FIRESS Instrument




- 4 slit-fed grating modules, each 24 x 84 pixels w/ gap.
- Bands 1 and 3 overlap, Bands 2 and 4 overlap.
- 2 pointings for full spectrum, though all 4 bands read out.
- High-res mode couples all bands when engaged

Simulated PRIMA/FIRESS spectra of two luminous galaxies at z > 4.5, with realistic noise model reflecting current best estimate FIRESS sensitivity. Dots are example PRIMAger binned fluxes. *From Donnely et al contribution to PRIMA GO book.*

PR:MA

FIRESS high-res FTM module






- When engaged, picks off light from telescope in collimated space
- Serves the full band simultaneously (2 pointings required for a source)
- Path length can be tuned, provides up to R=4,400 at 112 microns
- R scales as 1/lambda. So can resolve water lines in Band 1.

Heritage: Herschel SPIRE FTS also 4.5 K imaging FTS. (Griffin et al.) Same Canadian team developing the low-powerdissipation scan mechanism.




KID Detectors: a JPL / GSFC / SRON Collaboration for PRIMA


- Sensitivity exceeds performance requirements over full wavelength range.
- Demonstrated detector/lenslet hybridized arrays with full FIRESS format (84x12, 900-µm pixel pitch). PRIMAger prototypes in place from SRON.
- Key remaining challenge is yield with lenses bonded and full-band readout.



JPL FIRESS arrays 12 x 84 pixel tiles, 2 per focal plane: >90% resonator yield


///×/ ·9///×/>







PRIMA Sensitivities





Includes: all key optical efficiencies, microlens & pixel coupling to sources, grating blaze detector sensitivity, background photons temporal efficiencies due to chopping, cosmic ray removal.

PRIMA GO Opportunities

- Sensitivities available on fact sheet and web page.
 - PRIMAger (mapping speed)
 - FIRESS (pointed mapping #s on web page)
 - FIRESS high res (pointed)
- Instantaneous field of regard of 26% of the sky.
 - Investigating larger FoR current limits very conservative. Have margins in thermal lift and mass so could increase sunshade area.
- Agile. Can slew and anywhere in field of regard in 12 minutes
- No meaningful constraints on changing instruments or modes.
- GO book a set of example fiducial cases. Thanks very much to those of you that contributed. More to come.
- Remember, NASA will control the GO program.
- Reach out with questions.

prima.ipac.caltech.edu





Thank you

PRIMA activities at AAS (other than this session)

Monday 9-10: Poster Session 106: Steve Unwin et al., PRIMA Planetary System Formation

Tuesday 2-3:30: Oral 450: Margaret Meixner et al., PRIMA overview

Wednesday 9-10, Poster Session 306 Betsy Mills et al., PRIMA Galactic Ecosystems

Wednesday 5:30-6:30: Poster Session 306 Arielle Moullet et al.: PRIMA GO Book

Thursday 1-2, Poster Session 457 Matt Bradford et al. – FIRESS Margaret Meixner et al. – PRIMAger Johannes Staguhn et al. – Rise of Dust and Metals

Tuesday evening: Informal Social Event 8:30pm at the New Orleans Social House

PRIMA folks at AAS 243

.5



Lee Armus





Denis Burgarella





Tiffany Kataria

Margaret Meixner



Arielle Moullet

Steve Unwin

prima.ipac.caltech.edu

